

## 6 Waste (NIR Chapter 7)

For this methodology report, the Waste chapter consists of two subsectors: solid waste disposal and wastewater treatment and discharge. More information on national-level emissions and methods is available in Chapter 7 of the national *Inventory*. Note that emissions from waste incineration are discussed in Chapter 2, Section 2.1.4, of this methodology report. Table 6-1 summarizes the different approaches used to estimate state-level waste emissions and completeness across states. Geographic completeness is consistent with the national *Inventory*. The sections below provide more detail on each category.

**Table 6-1. Overview of Approaches for Estimating State-Level Waste Sector GHG Emissions and Sinks**

Category	Gas	Approach	Geographic Completeness <sup>a</sup>
Landfills	CH <sub>4</sub>	Approach 2	Includes emissions from industrial and municipal waste landfills from all states, the District of Columbia, tribal lands, and some territories (i.e., Guam, Puerto Rico) as applicable.
Wastewater	CH <sub>4</sub> , N <sub>2</sub> O	Approach 2	Includes emissions from all states, the District of Columbia, tribal lands, and some territories (i.e., American Samoa, Guam, Northern Mariana Islands, Puerto Rico, and U.S. Virgin Islands for domestic wastewater) as applicable.
Composting	CH <sub>4</sub> , N <sub>2</sub> O	Approach 2	Includes emissions from commercial composting facilities from all states, the District of Columbia, tribal lands, and some territories (Puerto Rico) as applicable.
Anaerobic Digestion at (Stand-Alone) Biogas Facilities	CH <sub>4</sub>	Approach 2	Includes emissions from all states, the District of Columbia, tribal lands, and territories.

<sup>a</sup> Emissions are likely occurring in other U.S. territories; however, due to a lack of available data and the nature of this category, this analysis includes emissions for only the territories indicated. Territories not listed are not estimated.

### 6.1 Solid Waste Disposal

This section presents the methodology used to estimate the emissions from solid waste disposal management activities, which consist of the following sources:

- Landfills (MSW and industrial waste) (CH<sub>4</sub>)
- Composting (CH<sub>4</sub>, N<sub>2</sub>O)
- Anaerobic digestion at biogas facilities (stand-alone) (CH<sub>4</sub>)

## 6.1.1 Landfills (NIR Section 7.1)

### 6.1.1.1 Background

After being placed in a landfill, organic waste such as paper, food scraps, and yard trimmings is initially decomposed by aerobic bacteria. After the oxygen has been depleted, the remaining waste is available for consumption by anaerobic bacteria, which break down organic matter into substances such as cellulose, amino acids, and sugars. These substances are further broken down through fermentation into gases and short-chain organic compounds that form the substrates for the growth of methanogenic bacteria. These CH<sub>4</sub>-producing anaerobic bacteria convert the fermentation products into stabilized organic materials and biogas consisting of approximately 50% biogenic CO<sub>2</sub> and 50% CH<sub>4</sub> by volume. CH<sub>4</sub> and CO<sub>2</sub> are the primary constituents of landfill gas generation and emissions. Consistent with the 2006 IPCC Guidelines, net CO<sub>2</sub> flux from C stock changes in landfills are estimated and reported under the LULUCF sector (see Section 3.4 of this report) (IPCC 2006).

More information on emission pathways and national-level emissions from landfills and associated methods can be found in the Waste chapter (Chapter 7), Section 7.1, of the national *Inventory* (EPA 2024), available online at [https://www.epa.gov/system/files/documents/2024-04/us-ghg-inventory-2024-chapter-7-waste\\_04-17-2024.pdf](https://www.epa.gov/system/files/documents/2024-04/us-ghg-inventory-2024-chapter-7-waste_04-17-2024.pdf).

### 6.1.1.2 Methods/Approach (Municipal Solid Waste Landfills)

The MSW landfill state emissions inventories apply Approach 2 for disaggregating national estimates and rely heavily on the Subpart HH data collected through the GHGRP. As explained in the methodology discussion of Section 7.1 of the national *Inventory*, EPA uses an IPCC Tier 2 approach and several data sources, methods, and assumptions to estimate emissions (see pages 7-8 through 7-9 for details on the inputs and equations). The state inventories apply a state percentage of either waste landfilled or net CH<sub>4</sub> emissions by state as reported to Subpart HH (EPA 2022) as a proxy for each state's share of CH<sub>4</sub> net emissions over the time series. Table 6-2 summarizes the methodology used to develop the state-level estimates, followed by additional detail. The annual state percentages were applied to the national estimates to retain an IPCC Tier 2 approach consistent with the national *Inventory*.

**Table 6-2. Summary of Approaches to Disaggregate the National *Inventory* for MSW Landfills Across Time Series**

Time Series Range	Summary of Method
1990–2009	<ul style="list-style-type: none"> <li>Applied the percentage of waste landfilled by state (aggregated total as reported by landfills in each state to Subpart HH for historical years) to the national CH<sub>4</sub> net emissions for each year (IPCC 2006 Tier 2)</li> <li>The state percentage approach accounts for all emissions, including those calculated in the national <i>Inventory</i> through back-casting Subpart HH data and scaling up emissions to account for smaller landfills that do not report through Subpart HH.</li> </ul>
2010–2022	<ul style="list-style-type: none"> <li>Applied the percentage of net CH<sub>4</sub> emissions by state (aggregated total as reported by landfills in each state to Subpart HH) to the national CH<sub>4</sub> net emissions for each year.</li> <li>The state percentage approach accounts for all emissions, including those calculated by scaling up emissions to account for smaller landfills that do not report through Subpart HH.</li> </ul>

Historical waste disposed of since a facility began operating is reported using prescribed methods in the rule to maintain consistency across the facility data. The quantity of waste landfilled by Subpart HH reporters was assumed to be representative of the universe of MSW landfills in the United States because Subpart HH reporters include each state's highest emitting MSW landfills, which is directly tied to the quantity of waste landfilled. The national *Inventory* methodology back-casts Subpart HH net CH<sub>4</sub> emissions and uses a scale-up factor to account for lower-emitting MSW landfills (e.g., non-reporters). The intent of the scale-up factor is to estimate CH<sub>4</sub> emissions from MSW landfills that do not report to the GHGRP. EPA has put significant effort into identifying landfills that do not report to the GHGRP, most recently in 2021. Basic landfill characteristics such as the landfill's name and location, first year of operation, current operational status, and waste-in-place data have been compiled for these landfills when available. Disaggregating the Subpart HH data by state was determined to be a reasonable assumption considering the lack of historical data for landfills that do not report to the GHGRP.

The methodology used for 1990–2009 applies a state percentage of waste landfilled for this time frame as reported by landfills under Subpart HH of the GHGRP to the national estimates of CH<sub>4</sub> emissions. Approximately 1,200 MSW landfills have reported to the GHGRP since reporting began in 2010. This approach disaggregates national net emissions values by applying the state percentage as a proxy of net emissions.

The methodology for 2010–2022 applies a state percentage of net CH<sub>4</sub> emissions reported by landfills under Subpart HH to the national estimates of CH<sub>4</sub> emissions. Using net CH<sub>4</sub> emissions is consistent with the recent methodological refinements in the national *Inventory* to incorporate reported Subpart HH net CH<sub>4</sub> emissions. Unlike the national *Inventory*, scale-up factors for each state were not developed since these would require significant effort; instead, the national emissions values are disaggregated by a proxy that is assumed to be generally representative of state-by-state emissions.

Emissions from managed landfills located in Puerto Rico and Guam are included because facilities in these territories report to Subpart HH.

### 6.1.1.3 Methods/Approach (Industrial Landfills)

The state inventories estimate CH<sub>4</sub> emissions from industrial waste landfills for two industry categories consistent with the national *Inventory*: (1) pulp and paper and (2) food and beverage. Data reported to the GHGRP on industrial waste landfills suggest that most of the organic waste that would result in CH<sub>4</sub> emissions is disposed of at pulp and paper and food processing facilities. Information on both industry categories with respect to the amount of waste generated and disposed of in a dedicated industrial waste landfill is limited; thus, EPA uses a Tier 1 approach to estimate CH<sub>4</sub> emissions. Additionally, no comprehensive list of industrial waste landfills exists. While the information is available in the *Waste Business Journal* (WBJ), the date of data related to each waste management facility included is unknown. Therefore, EPA does not have information on the number of industrial waste landfills that were operational over the time series and information regarding the number of industrial waste landfills located in each state. The types and amounts of waste disposed of in the operational industrial waste landfills are also limited.

A portion of pulp and paper mills in the United States report to Subpart TT (Industrial Waste Landfills) of the GHGRP. Previous analyses of the 2016 pulp and paper emissions from the GHGRP (RTI International 2018) showed that total Subpart TT emissions from facilities associated with a pulp and paper NAICS code generally align (within approximately 10–20%) with the national *Inventory*'s national estimate of emissions from the pulp and paper manufacturing sector. On the other hand, a small number of facilities associated with a food and beverage NAICS code report to Subpart TT, and these emissions are vastly different between Subpart TT and the national *Inventory*.

Because of the data limitations described above, Approach 2 was used to disaggregate the national *Inventory* CH<sub>4</sub> emissions for both industry categories, rather than a more detailed facility-specific, bottom-up approach.

#### **6.1.1.3.1. Pulp and Paper Manufacturing**

For the pulp and paper source category, EPA extracted a state-by-state count of mills in the United States from two sources: Data Basin for 2008 and Mills OnLine for 2015–2016 (Conservation Biology Institute 2008; Ga Tech Center for Paper Business and Industry Studies n.d.). The count of facilities is approximately 233 and 332 from Data Basin and Mills OnLine, respectively. The count and percentage of mills by state are shown in Appendix F (Table F-1). According to the Industrial Resources Council, mills are located in 41 states, not including Alaska, Colorado, North Dakota, Nebraska, Nevada, Rhode Island, South Dakota, Utah, and Wyoming. For comparison, the Subpart TT pulp and paper facilities across RYs 2011–2019 represent a maximum of 92 facilities located across 21 states.

To estimate CH<sub>4</sub> generation and emissions, the Data Basin 2008 percentages by state were applied to the national *Inventory* estimate for the pulp and paper manufacturing sector for 1990–2010, and the Mills OnLine 2015–2016 percentages by state were applied for 2011–2022. This approach assumes broadly that each facility is generating an equal amount of waste that is landfilled and, therefore, an equal amount of CH<sub>4</sub> emissions. Consistent with the national *Inventory*, this assumption and this approach were used in an attempt to ensure complete coverage of industrial waste landfills in the United States because the Subpart TT pulp and paper facilities may not equal the total number of pulp and paper facilities disposing of waste in dedicated industrial waste landfills. The exact number of pulp and paper manufacturing facilities that dispose of waste in industrial waste landfills is unknown.

CH<sub>4</sub> emissions from the pulp and paper sector were disaggregated by applying the percentage of the mills by state as a proxy for facilities generating and disposing of waste in industrial waste landfills. No additional calculations were performed, and the IPCC Tier 1 methodology (IPCC 2006) used to generate the national emissions estimates was applied by default.

#### **6.1.1.3.2. Food and Beverage Manufacturing**

Minimal data are available to characterize the amounts and types of waste generated nationally from food and beverage manufacturers and disposed of in industrial waste landfills. Less is known about the number of facilities in each state that dispose of waste in a dedicated industrial landfill.

A similar approach using a count of assumed industrial food and beverage manufacturing facilities that dispose of waste in an industrial waste landfill by state was applied to the national food and beverage category estimates. The list of food and beverage manufacturing facilities consists of 13 NAICS codes as shown in Appendix F (Table F-2) comprising 9,175 facilities. This list (which can be shared on request) was extracted from the 2021 update to the EPA Excess Food Opportunities database (EPA 2021).

The EPA Excess Food Opportunities database includes a low- and high-end estimate of the amount of excess food generated (tons/year). These data were not used in the methodology. Rather, the average percentage of the amount of excess food generated by each state across the selected NAICS codes was used as a proxy for the share of CH<sub>4</sub> generation and emissions estimates. The same approach used for the pulp and paper manufacturing sector was applied whereby the average percentage of excess food by state was applied to the national total amount of CH<sub>4</sub> generation and CH<sub>4</sub> emissions for each year of the time series. This is a broad assumption but allows for the calculation of emissions with limited knowledge on the locations of facilities disposing of food waste into industrial waste landfills.

The percentage of excess food generated by state is presented in Appendix F (see Table F-3). Note that the Excess Food Opportunities database and map do not indicate the management pathway for the excess food. The EPA Facts and Figures methodology (EPA 2020) also does not include an estimation of the amount of excess food being disposed of in industrial waste landfills. Therefore, the percentage of waste disposed of is likely overrepresented for some states and is why the estimates for the District of Columbia, the Virgin Islands, and Puerto Rico have been zeroed out.

#### 6.1.1.4 Recalculations

No recalculations were applied for this current report.

#### 6.1.1.5 Uncertainty

The overall uncertainty associated with the 2022 national estimates of CH<sub>4</sub> from MSW and industrial waste landfills was calculated using the Approach 2 methodology (IPCC 2006). As described further in Chapter 7 of the national *Inventory*, levels of uncertainty in the national estimates in 2022 were -2%/+20% of the estimated CH<sub>4</sub> emissions for MSW landfills and -31%/+25% for industrial waste landfills.

State-level estimates likely have a higher uncertainty due to (1) apportioning the national emissions estimates to each state based on assumptions made to disaggregate the national emissions estimates, which are based on state percentages as reported to the GHGRP, and (2) the application of the scale-up factor to nationally compiled landfill gas recovery databases used in the national *Inventory*. Additionally, state-level estimates before the GHGRP began (i.e., before 2010) may have more uncertainty than state-level estimates after the GHGRP began (i.e., 2010 and afterward). For more details on national level uncertainty, see the uncertainty discussion in Section 7.1 of the national *Inventory*.

#### 6.1.1.6 Planned Improvements

Potential refinements to landfill estimation methods include the following:

- **MSW landfills.** Planned improvements to the state-level estimates are consistent with those presented in Section 7.1 of the national *Inventory*. In particular, EPA plans to improve completeness of emissions from all waste management practices (i.e., open dumpsites) in U.S. territories by identifying data and applying methods to include emissions from open dumpsites in territories.
- **Industrial waste landfills.** A more complete and comprehensive list of pulp and paper facilities in the United States will be identified, including years of operation since 1990. Further QC on this inventory will be performed by comparing the counts of industrial waste landfills by state in available data sets.

#### 6.1.1.7 References

Center for Paper Business and Industry Studies (n.d.) *Mills Online 2015-2016*. Available online at: [https://epay.gatech.edu/C20793\\_ustores/web/product\\_detail.jsp?PRODUCTID=2466](https://epay.gatech.edu/C20793_ustores/web/product_detail.jsp?PRODUCTID=2466).

Conservation Biology Institute (2008) *Data Basin: U.S. Pulp and Paper Mills*. Available online at: <https://databasin.org/maps/new/#datasets=1f2a22ee1aa441568cbf5bea1b275c88>.

EPA (U.S. Environmental Protection Agency) (2020) *Advancing Sustainable Materials Management: 2018 Tables and Figures*. Available online at: [https://www.epa.gov/sites/production/files/2020-11/documents/2018\\_tables\\_and\\_figures\\_fnl\\_508.pdf](https://www.epa.gov/sites/production/files/2020-11/documents/2018_tables_and_figures_fnl_508.pdf).

EPA (2021) Excess Food Opportunities Map, Available online at: <https://www.epa.gov/sustainable-management-food/excess-food-opportunities-map>.

EPA (2022) *Envirofacts Data. Subpart HH: Municipal Solid Waste Landfills*.

EPA (2024) *Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990–2022*. EPA 430-R-24-004. Available online at: <https://www.epa.gov/ghgemissions/inventory-us-greenhouse-gas-emissions-and-sinks>.

IPCC (Intergovernmental Panel on Climate Change) (2006) *2006 IPCC Guidelines for National Greenhouse Gas Inventories*. H.S. Eggleston, L. Buendia, K. Miwa, T. Ngara, and K. Tanabe (eds.). Institute for Global Environmental Strategies. Available online at: <https://www.ipcc-nggip.iges.or.jp/public/2006gl/>.

RTI International (2018) *Comparison of Industrial Waste Data Reported Under Subpart TT and the Solid Waste chapter of the GHG Inventory*. Memorandum prepared by K. Bronstein, B. Jackson, and M. McGrath for R. Schmeltz (EPA).

## 6.1.2 Composting (NIR Section 7.3)

### 6.1.2.1 Background

This section presents methods used to estimate state-level GHGs from large-scale commercial composting facilities that typically include sections of the waste that operate in an anaerobic environment where degradable organic carbon in the waste material is broken down, generating CH<sub>4</sub> and N<sub>2</sub>O. Even though CO<sub>2</sub> emissions are generated, they are not included in net GHG emissions for composting. Consistent with the national *Inventory*, emissions from residential (backyard) composting are not included in the scope. Additionally, the national *Inventory* assumes windrow is the composting method used, and the waste mixture is homogeneous, consisting primarily of yard waste and some food. Annual throughput data on static and in-vessel commercial composting methods were not identified in secondary (published) data. Consistent with the 2006 IPCC Guidelines, net CO<sub>2</sub> flux from C stock changes in waste material is estimated and reported under the LULUCF sector (see Chapter 3.4 of this report) (IPCC 2006).

More information on emission pathways and national-level emissions from composting and associated methods can be found in the Waste chapter (Chapter 7), Section 7.3 of the national *Inventory*, available online at [https://www.epa.gov/system/files/documents/2024-04/us-ghg-inventory-2024-chapter-7-waste\\_04-17-2024.pdf](https://www.epa.gov/system/files/documents/2024-04/us-ghg-inventory-2024-chapter-7-waste_04-17-2024.pdf).

### 6.1.2.2 Methods/Approach

EPA compiles national CH<sub>4</sub> and N<sub>2</sub>O emissions estimates for commercial composting facilities in the United States using an IPCC Tier 1 method by which an IPCC default emissions factor is applied to the national quantity of material composted. No facility-specific information is used because it is generally unavailable over the time series.

The national *Inventory* was disaggregated to the state level using Approach 2 on the basis of data available for the proportion of material composted by state for select years. Table 6-3 summarizes published state-level estimates of composted material used in this inventory. Years where published data are not available are either interpolated or extrapolated using population growth and published estimates.

**Table 6-3. Summary of Availability and Sources for Composting Data**

Year Composting Data Available for	Reference Citation
2000	Goldstein and Madtes 2001
2002	Kaufman et al. 2004
2004	Goldstein et al. 2006
2006	Arsova et al. 2008
2008	Arsova et al. 2010
2010	EREF 2016
2011	Shin 2014
2012	Platt 2014
2013	EREF 2016
2016	WBJ 2016
2020	WBJ 2020

The state-level data were largely compiled from voluntary surveys of state agencies that reported MSW generated and estimates by relevant management pathways (e.g., landfill, recycling, composting). Composting estimates may be directly reported by the state agencies or estimated or adjusted by the report authors using the best available information for available years. Occasionally, data for some states are not available and are indicated as such in the data sources. The WBJ is an annually updated database of which the quality is unknown, but it is used because there is a general lack of data. Both the WBJ 2016 and 2020 were used to estimate state data for 2017–2019. Completeness is one limitation with the available state data used.

The general methodology to estimate the annual quantity of waste composted per year is as follows:

$$Composted_s = \%_s \times N_C$$

where:

$Composted_s$	= the mass of material composted by state (tons/year)
$\%_s$	= the state percentage of material composted, calculated using available state data (%)
$N_C$	= the national estimate of material composted as reported in the EPA Advancing Sustainable Materials Facts and Figures reports (tons/year) (EPA 2020)

The state percentages of material composted were calculated by dividing each state-reported amount of waste composted by the total of all material composted for that year. The sum of all state-reported data is referred to as national estimates by the report authors, but to avoid confusion with the Facts and Figures data published by EPA, this methodology report refers to this as “the sum of state-reported data.”. Limitations with the state-reported survey data include its voluntary nature and the occasional lack of data for states that did not provide a survey response. The report authors noted they made assumptions to estimate and adjust data to the extent possible. For years where no state data were reported in a specific survey, EPA estimated the data using the prior or next year of available data. These gaps were minimal (i.e., five or fewer states for each survey year).

Because state data are only available for select years, interpolation and extrapolation were required to generate estimates for each year of the time series. State proportions applied to 1990–1999 are the same as those for 2000 (Goldstein and Madtes 2001). No state data exist for this portion of the time series, and there

is a large amount of uncertainty surrounding the number of facilities and amount of material composted. This is a conservative approach since it is unknown when a state began compositing operations, so it is assumed if they had operations in 2000 that they did in 1990 as well. Data in between the survey data were interpolated using the prior year's and next year's survey data (the state proportion of material composted). Annual state data were interpolated for 2001, 2003, 2005, 2007, 2009, 2014, 2017, 2018, and 2019. Annual state data for 2021 and 2022 were extrapolated using population growth (U.S. Census Bureau 2022) and estimates of material composted (WBJ 2020). State percentages for each year are presented in Appendix F (Table F-4).

The formula used for interpolation of the state percentage for the year in question is as follows:

$$y = \left( \frac{y_2 - y_1}{x_2 - x_1} \right) \times (x) - x_1 + y_1$$

where:

$y$	= state percentage of waste composted for the year without data, %
$y_1$	= state percentage of waste composted for the prior year with data, %
$y_2$	= state percentage of waste composted for the next year with data, %
$x$	= the year without data
$x_1$	= the prior year with data
$x_2$	= the next year with data

The state percentage data were multiplied by the national estimate of material composted from the EPA Facts and Figures reports to cap the total quantity composted across the states and match the state totals to the national *Inventory*. The EPA Facts and Figures national estimates were directly used to estimate the national *Inventory*. The IPCC Tier 1 method used in the national *Inventory* estimates (IPCC 2006) is the product of an emissions factor and the mass of organic waste composted.

The final step in developing the state inventory was estimating the CH<sub>4</sub> and N<sub>2</sub>O emissions. For simplicity, the state percentages were multiplied by the annual national emissions estimates.

### 6.1.2.3 Recalculations

No recalculations were applied for this current report.

### 6.1.2.4 Uncertainty

The overall uncertainty associated with the 2022 national estimates of CH<sub>4</sub> and N<sub>2</sub>O from composting (specifically large-scale, commercial composting facilities) was calculated using the 2006 IPCC Guidelines Approach 1 methodology (IPCC 2006). As described further in Chapter 7 of the national *Inventory*, levels of uncertainty in the national estimates in 2022 were -58%/+58% for CH<sub>4</sub> and for N<sub>2</sub>O. State-level estimates will have a higher uncertainty than the national estimates because of apportioning the national quantity of material composted (sourced from the EPA Sustainable Materials Management reports and calculated with a mass balance methodology) to each state based on sporadically published waste management data from a voluntary state agency survey for select years. The national methodology also assumes most composting in the United States uses the windrow method and treats a homogeneous mixture of primarily yard trimmings and some food waste. For more details on national-level uncertainty, see the uncertainty discussion in Section 7.3 of the national *Inventory*, available online at [https://www.epa.gov/system/files/documents/2024-04/us-ghg-inventory-2024-chapter-7-waste\\_04-17-2024.pdf](https://www.epa.gov/system/files/documents/2024-04/us-ghg-inventory-2024-chapter-7-waste_04-17-2024.pdf).

### 6.1.2.5 Planned Improvements

In future annual publications, EPA plans to investigate state volumes of material composted where the report authors (from referenced composting data sources) indicated potentially combined volumes of waste



sent to composting, recycling, and anaerobic digestion. EPA will continue to identify annual quantities of material composted in states where data are lacking (e.g., Alaska, Guam). For example, a 2021 desk-based investigation into composting facilities in Alaska revealed operational aerated composting facilities, but the annual capacity and throughput were not identified. EPA will continue to search for relevant data for commercial composting facilities in these states. Planned improvements to the national estimates for composting outlined in Section 7.3 (page 7-58) of the national *Inventory* will lead directly to improvements in the quality of state-level estimates as well.

#### 6.1.2.6 References

- Arsova, L., R. van Haaren, N. Goldstein, S.M. Kaufman, and N.J. Themelis (2008) The State of Garbage in America. *BioCycle*. Available online at: <https://www.biocycle.net/the-state-of-garbage-in-america-3/>.
- Arsova, L., R. Van Haaren, N. Goldstein, S. Kaufman, and N. Themelis (2010) The State of Garbage in America. *BioCycle*, 51(10): 16. Available online at: <https://www.biocycle.net/2010/10/26/the-State-of-garbage-in-america-4/>.
- EPA (U.S. Environmental Protection Agency) (2020) *Advancing Sustainable Materials Management: 2018 Tables and Figures*. Available online at: [https://www.epa.gov/sites/production/files/2020-11/documents/2018\\_tables\\_and\\_figures\\_fnl\\_508.pdf](https://www.epa.gov/sites/production/files/2020-11/documents/2018_tables_and_figures_fnl_508.pdf).
- EREF (Environmental Research & Education Foundation) (2016) *Municipal Solid Waste Management in the United States: 2010 & 2013*.
- Goldstein, N., and C. Madtes (2001) The State of Garbage in America. *BioCycle*.
- Goldstein, N., S. Kaufman, N. Themelis, and J. Thompson Jr. (2006) The State of Garbage in America. *BioCycle*, 47(4): 26. Available online at: <https://www.biocycle.net/2006/04/21/the-state-of-garbage-in-america-2/>.
- IPCC (Intergovernmental Panel on Climate Change) (2006) *2006 IPCC Guidelines for National Greenhouse Gas Inventories*. H.S. Eggleston, L. Buendia, K. Miwa, T. Ngara, and K. Tanabe (eds.). Institute for Global Environmental Strategies. Available online at: <https://www.ipcc-nggip.iges.or.jp/public/2006gl/>.
- Kaufman, S.M., N. Goldstein, K. Millrath, and N.J. Themelis (2004) The State of Garbage in America. *BioCycle*, 45(1): 31. Available online at: <https://www.biocycle.net/the-state-of-garbage-in-america/>.
- Platt, B. (2014) *State of Composting in the US: What, Why, Where & How*. Institute for Local Self-Reliance. Available online at: <https://ilsr.org/articles/state-of-composting/>.
- Shin, D. (2014) *Generation and Disposition of Municipal Solid Waste (MSW) in the United States—A National Survey*. Thesis. Columbia University, Department of Earth and Environmental Engineering, January 3, 2014.
- U.S. Census Bureau (2022) Table NST-EST2022-POP. In: *Annual Estimates of the Resident Population for the United States, Regions, States, District of Columbia, and Puerto Rico: April 1, 2020 to July 1, 2022*. Release date: December 2022.
- WBJ (Waste Business Journal) (2016) *Directory of Waste Processing & Disposal Sites 2016*.
- WBJ (2020) *Directory of Waste Processing & Disposal Sites 2020*.

## 6.1.3 Anaerobic Digestion at Biogas Facilities (Stand-Alone) (NIR Section 7.4)

### 6.1.3.1 Background

Anaerobic digestion is a series of biological processes in the absence of oxygen in which microorganisms break down organic matter, producing biogas and soil. Stand-alone digestion is one of three main categories of anaerobic digestion facilities, which also includes on-farm digesters and digesters at water resource recovery facilities. This section focuses exclusively on stand-alone digesters, which typically manage food waste from different sources, including food and beverage processing industries. Emissions from on-farm digesters and digesters at water resource recovery facilities are reflected under Sections 4.1.2 (Manure Management) and 6.2.1 (Wastewater Treatment and Discharge) of this report. Based on available data, anaerobic digestion of food waste occurs in 31 states, listed in Appendix F (Table F-5).

At stand-alone digestors, CH<sub>4</sub> emissions may result from a fraction of the biogas lost during the process due to leakages and other unexpected events (0–10% of the amount of CH<sub>4</sub> generated; IPCC 2006). The remaining biogas (90–100% of gas generated) is flared or used beneficially, often combusted to produce heat and power, or further processed into renewable natural gas or for use as a transportation fuel. CO<sub>2</sub> emissions are biogenic in origin and should be reported as an informational item in the energy sector (IPCC 2006).

More information on emission pathways and national-level emissions and methods can be found in Section 7.4 of the national *Inventory*.

### 6.1.3.2 Methods/Approach

EPA compiles national CH<sub>4</sub> emissions estimates for stand-alone anaerobic digester facilities in the United States using an IPCC Tier 1 method, which applies an IPCC default leakage factor of 5% to the CH<sub>4</sub> generated. The amount of CH<sub>4</sub> generated is the product of an emission factor and the mass of organic waste processed. The weighted average annual quantity of material processed is estimated from voluntary EPA AD Survey data (EPA 2018, 2019, 2021, 2023) and an estimated number of operating facilities per year (see Table 7-46 and Table 7-47, respectively, of the national *Inventory*). No facility-specific quantities of material digested were directly used because of a general lack of facility-specific data over the time series. The methodology applied to generate the national *Inventory* was based on two large assumptions—the number of operational facilities and the weighted average of material digested for two of the 30 years in the time series (1990–2022). The state inventory further takes these assumptions to a state level by assuming that the same percentage of total operational facilities is the same for each year of the time series because of a general lack of data on total operational facilities by state across the time series. Therefore, the state-level inventories are a gross estimate that may be refined in future years if available information by state is obtained.

In the national *Inventory*, EPA calculated a weighted average of material digested using masked survey data from available survey reports for 2015 to 2019 (EPA 2018, 2019, 2021, 2023). The weighted average was applied to an estimated number of operational facilities per year to estimate the annual quantity of material digested. The first step to calculating the state inventory was to disaggregate the annual estimates of the material digested. This was disaggregated by applying a state percentage of operational facilities as reported to the two published EPA survey reports (EPA 2018, 2019). The state proportions of operational facilities in 2015 and 2016 are presented in Appendix F (Table F-5).

The state proportions were multiplied by the national quantity digested for each year in the time series, which forced the total quantities across the states to match the national *Inventory* estimates. The same state percentage was used for each year in the time series because of a lack of compiled data on the number of stand-alone digesters by state between 1990 and 2022.

### 6.1.3.3 Recalculations

For the current *Inventory*, a methodological change was made whereby the CH<sub>4</sub> emissions are considered equal to leakage from the digester network of pipes. A leakage factor of 5% (as recommended in IPCC 2006) is applied to the CH<sub>4</sub> generation estimate for all years in the time series. This methodological change applies to every year in the time series and significantly reduces annual CH<sub>4</sub> emissions estimates. Previously, the EPA AD Survey data reporting the amount of biogas produced at AD facilities was used for the amount of gas recovered, with the remaining gas assumed to be leaked or emitted. This method calculated higher emissions estimates, which showed the majority of the gas generated at an AD being emitted instead of being used in biogas projects. This was inconsistent with the EPA AD Survey findings that approximately 95% of stand-alone AD facilities use some or all biogas on-site, and it was also inconsistent with the IPCC guidance on default leakage from AD facilities. EPA will further investigate the survey data for the total biogas-produced, since they indicate very low gas utilized as compared to this revised methodology. See the recalculations discussion (page 7-63) in Section 7.4 of the national *Inventory*.

### 6.1.3.4 Uncertainty

The overall uncertainty associated with the 2022 national estimates of CH<sub>4</sub> from stand-alone anaerobic digesters was calculated using the 2006 IPCC Guidelines Approach 1 methodology (IPCC 2006). As described further in Chapter 7 of the national *Inventory*, levels of uncertainty in the national estimates in 2019 were -54%/+54% CH<sub>4</sub>. State-level estimates will have a higher uncertainty because of apportioning the national emissions estimates to each state based solely on the number of stand-alone anaerobic digester facilities from EPA survey data collected between 2015 and 2018. Emissions estimates before 2015 will have a higher uncertainty than those in 2015 and later years. These assumptions were required because of limited facility-specific data presented in secondary resources. For more details on national level uncertainty, see the uncertainty discussion in Section 7.4 of the national *Inventory*.

### 6.1.3.5 Planned Improvements

The planned improvements are consistent with those planned for improving national estimates given that the underlying methods for state GHG estimates are the same as those in the national *Inventory*. To find information on planned improvements to refine methods for estimating emissions from stand-alone anaerobic digestion, see the planned improvements discussion starting on page 7-64 of Section 7.4 in the national *Inventory*.

### 6.1.3.6 References

- EPA (U.S. Environmental Protection Agency) (2018) *Anaerobic Digestion Facilities Processing Food Waste in the United States in 2015: Survey Results*. EPA/903/S-18/001. Available online at: [https://www.epa.gov/sites/default/files/2018-08/documents/ad\\_data\\_report\\_final\\_508\\_compliant\\_no\\_password.pdf](https://www.epa.gov/sites/default/files/2018-08/documents/ad_data_report_final_508_compliant_no_password.pdf).
- EPA (2019) *Anaerobic Digestion Facilities Processing Food Waste in the United States in 2016: Survey Results*. EPA/903/S-19/001. Available online at: [https://www.epa.gov/sites/default/files/2019-09/documents/ad\\_data\\_report\\_v10\\_-\\_508\\_comp\\_v1.pdf](https://www.epa.gov/sites/default/files/2019-09/documents/ad_data_report_v10_-_508_comp_v1.pdf).
- EPA (2021) *Anaerobic Digestion Facilities Processing Food Waste in the United States (2017 & 2018): Survey Results*. EPA/903/S-21/001. Available online at: [https://www.epa.gov/sites/default/files/2021-02/documents/2021\\_final\\_ad\\_report\\_feb\\_2\\_with\\_links.pdf](https://www.epa.gov/sites/default/files/2021-02/documents/2021_final_ad_report_feb_2_with_links.pdf).
- EPA (2023) *Anaerobic Digestion Facilities Processing Food Waste in the United States (2019): Survey Results*. EPA 530-R-23-003. Available online at: [https://www.epa.gov/system/files/documents/2023-04/Anaerobic\\_Digestion\\_Facilities\\_Processing\\_Food\\_Waste\\_in\\_the\\_United\\_States\\_2019\\_20230404\\_508.pdf](https://www.epa.gov/system/files/documents/2023-04/Anaerobic_Digestion_Facilities_Processing_Food_Waste_in_the_United_States_2019_20230404_508.pdf).

IPCC (Intergovernmental Panel on Climate Change) (2006) *2006 IPCC Guidelines for National Greenhouse Gas Inventories*. H.S. Eggleston, L. Buendia, K. Miwa, T. Ngara, and K. Tanabe (eds.). Institute for Global Environmental Strategies. Available online at: <https://www.ipcc-nggip.iges.or.jp/public/2006gl/>.

## 6.2 Wastewater Management

This section presents the methodology used to estimate the emissions from domestic and industrial wastewater treatment and discharge (CH<sub>4</sub>, N<sub>2</sub>O).

### 6.2.1 Wastewater Treatment and Discharge (NIR Section 7.2)

#### 6.2.1.1 Background

Consistent with the national *Inventory* and international guidance, EPA has developed disaggregated state estimates for both domestic and industrial wastewater treatment and discharge, as discussed below:

- Domestic wastewater CH<sub>4</sub> and N<sub>2</sub>O emissions originate from both septic systems and centralized treatment plants. Within these centralized plants, CH<sub>4</sub> emissions can arise from aerobic systems that liberate dissolved CH<sub>4</sub> that formed within the collection system or that are (1) designed to have periods of anaerobic activity, (2) from anaerobic systems, and (3) from anaerobic sludge digesters when the captured biogas is not completely combusted. N<sub>2</sub>O emissions can result from aerobic systems as a byproduct of nitrification, or as an intermediate product of denitrification. Methane emissions will also result from the discharge of treated effluent from centralized treatment plants to water bodies where carbon accumulates in sediments, while N<sub>2</sub>O emissions will also result from discharge of centrally treated wastewater to water bodies with nutrient-impacted or eutrophic conditions.
- Industrial wastewater CH<sub>4</sub> emissions originate from in-plant treatment systems, typically comprising biological treatment operations in which some operations are designed to have anaerobic activity or may periodically form anaerobic conditions. N<sub>2</sub>O emissions are primarily expected to occur from aerobic treatment systems as a byproduct of nitrification, or as an intermediate product of denitrification. Emissions will also result from discharge of treated effluent to waterbodies.

#### 6.2.1.2 Methods/Approach (Domestic Wastewater)

EPA estimated state-level domestic wastewater treatment and discharge emissions (CH<sub>4</sub>) using a simplified approach to apportion the national emission estimates to each state based on population (i.e., Approach 2 as defined in the Introduction to this report) and state-level septic data. In this method, EPA accessed historical U.S. Census data to compile state-level population data for each year of the inventory (1990–1999: U.S. Census Bureau 2002; 2000–2009: U.S. Census Bureau 2011; 2010–2021: U.S. Census Bureau 2021a, 2021b, 2022, 2023; Instituto de Estadísticas de Puerto Rico 2021). The U.S. Census Bureau (1990) and NEBRA (2022) reported the percentage of the population associated with septic systems in each state for 1990 and 2018, respectively. These percentages were multiplied by the 1990 and 2018 state-level population and then divided by the total summed national population to estimate the percentage of the national population with a septic system in each state and territory in 1990 and 2018. The state-level percentages for 1991–2017 were linearly interpolated between 1990 and 2018, and the remainder of the time series was set equal to 2018, as shown in Appendix F, Table F-6.

EPA calculated state- and territory-level emissions by multiplying the proportion of the U.S. population on centralized treatment or septic systems in each state or territory by the national CH<sub>4</sub> and N<sub>2</sub>O emissions for each year of the time series.

This simplified approach assumes the following:

- Every state has the same wastewater treatment system usage as the national *Inventory*.
- Every state has same distribution of discharge to various waterbody types as the national *Inventory*.
- Kitchen disposal usage is the same in every state, and wastewater biochemical oxygen demand (BOD) produced per capita, with and without kitchen scraps, is the same in every state (i.e., assumes total wastewater BOD produced per capita is the same as national production).
- Per capita protein consumption in the United States is the same in every state (i.e., assumes per capita consumption is the same as national consumption).

EPA did not perform a more detailed approach that would account for the specific types of treatment at centralized systems, such as anaerobic reactors or activated sludge, used in each state (see planned improvements below in Section 6.2.1.6). Similarly, there are insufficient readily available data sources to allow classification of the type of specific water bodies within each state, so EPA did not consider the type of water body receiving wastewater discharges within each state.

### 6.2.1.3 Methods/Approach (Industrial Wastewater)

Consistent with the national *Inventory* and national estimates, both CH<sub>4</sub> and N<sub>2</sub>O emissions were estimated for treating industrial wastewater from pulp and paper manufacturing, meat and poultry processing, petroleum refining, and breweries, while CH<sub>4</sub> emissions were also estimated for treating industrial wastewater from vegetables, fruits, and juices processing, and for starch-based ethanol production. These are the industry categories that are likely to produce significant GHG emissions from wastewater treatment. Data on industrial production by state are available or can be estimated from other readily available data for at least some of the time series of the inventory.

EPA estimated state-level emissions by estimating the percentage of the industry production that occurs in each state (i.e., using Approach 2 as described in the Introduction to this report). Where data were readily available, EPA estimated the distribution of production for each year of the time series and multiplied that by the national emissions estimate for each year of the time series. In some cases, due to time and resources, EPA was able to estimate the distribution of production for a subset of years in the time series, as discussed below by industry.

For pulp and paper manufacturing, state-level production data are not available, so EPA estimated state-level emissions by estimating the percentage of wastewater directly discharged in that state compared to the total flow of wastewater directly discharged for that industry, using data reported to EPA's ICIS National Pollutant Discharge Elimination System (NPDES) database. EPA acknowledges that this methodology ignores production at mills that either do not discharge wastewater or that discharge to a publicly owned treatment works. In both cases, these mills could be performing on-site treatment and emitting GHGs that cannot be captured.

EPA then multiplied that percentage by the national emissions estimate to obtain a state-level emissions estimate. Because of the limitation of data resources for this effort, EPA accepted most ICIS-NPDES data as is, but some outliers were determined and handled as described below (see planned improvements below in Section 6.2.1.6).

Both approaches assume the following:

- All facilities in an industry within a state have the same distribution of wastewater treatment operations as the national distribution.

- Every state has the same BOD and total nitrogen in untreated industry wastewater as the national-level estimates.
- Every state has the same nitrogen removal factor as the national-level estimates.
- The percentage of wastewater directly discharged by the state represents the distribution of all pulp and paper production by the state.

Further details on methods and data sources assumptions for each industry treating wastewater are described below.

#### 6.2.1.3.1. *Pulp and Paper Manufacturing*

- Industrial production data for pulp and paper are highly confidential and are not available by state.
- EPA used the amount of wastewater directly discharged by pulp mills by state—reported to both ICIS- NPDES from Enforcement Compliance History Online (EPA 2024b) and the Washington Department of Ecology’s Permitting and Reporting Information System, or PARIS (Washington Department of Ecology 2022, 2024)—to proportion U.S. national emissions estimates to a state (as shown in Appendix F, Table F-7). Because wastewater flow data housed in ECHO changed in 2016, using older data may cause discontinuities in the time series. EPA determined the distribution of discharge flow by state for 2019–2022 using ECHO and PARIS data and applied the 2019 distribution to all prior years of the national *Inventory*. There was no wastewater flow reported for the District of Columbia or U.S. territories for this industry.
  - Pulp and paper mills were determined in ECHO using Standard Industrial Classification codes 2611, 2621, and 2631.
- For facilities in states other than Washington, EPA:
  - Downloaded the total pulp and paper permit universe in ECHO, including permits that have discharge monitoring report data (252 facilities in 2022), and permits with information only (e.g., facility address) (322 facilities in 2022).
    - Stormwater, construction, or non-mill-related permits that were reported for a facility that also reported using another permit (such as a major or general permit) were removed from the analysis to prevent overestimating flow (see Table 1). If a facility only reported using stormwater permits, a single stormwater permit was retained so that it would be counted in the universe and flow could be estimated for the facility.
    - In four cases, it was discovered that a permit for an operational facility was missing from the data download for certain years, so those permit numbers were manually added to the facility universe for the missing years and the flow data for those facilities were estimated following the methodology (see Table 1).
  - Downloaded 2019–2022 flow data where available (EPA 2024b). Not all facilities report total flow if it is not required by their permit. Total flow was summed by state.
    - EPA determined four state flow outliers, one for Missouri in 2020, West Virginia in 2021 and 2022, and one for Minnesota in 2022. Outliers, determined as values that are at least an order of magnitude larger (10 times) than other years’ values for the state, were removed. It is assumed these values are data entry errors in ECHO. An average of the other available values was used as a surrogate for removed values.
  - For permits without flow data, total flow was estimated by using average flow by state, or average national total flow for that year if no state data were available, multiplied by the number of permits without flow data for that state.

- Facilities located in the state of Washington are not currently reported within ECHO due to lagging electronic reporting. To fill this known gap, EPA investigated a separate source for these data and:
  - Downloaded and reviewed permit data for known pulp mills determined from the Washington Department of Ecology’s Industrial Facility Permits website.
  - Downloaded 2019 flow data where available (Washington Department of Ecology 2022) for monitoring locations that are associated with process wastewater, per the facility permit. During the 1990–2022 *Inventory* update cycle, a new Washington permit that had since become active (WA0002925) was added to the 2022 estimate (Washington Department of Ecology 2024).
  - Multiplied the daily flow rate by 365.25 days to estimate a total yearly flow, then multiplied by number of months data were reported (to prevent overestimating annual flow, which was done to better match the methodology in ECHO).
  - Integrated into the other state data for all years.
- EPA calculated the percentage of national flow by state:
  - As with Washington, some states are missing from ECHO (e.g., Montana, Colorado). EPA assumed some of these states have nonzero emissions, but they do not have the data to determine whether there are facilities present or to estimate emissions, so they are reported as not applicable.
- EPA calculated the state-level emissions by multiplying national emissions by the percentage of national flow by state.
- Example: 2022 Georgia emissions
  - Georgia has 18 facilities in the facility universe, of which 14 have reported annual flow data.
  - The total flow based on the sum of reported flows (14 facilities) and calculated flows (4 facilities) from the state average flow of 8,042 million gallons (MMGal) for all facilities was 144,760 MMGal in 2022.
  - Georgia’s flow was 8.59% of the total national total flow of (1,684,263 MMGal).
  - Pulp and paper’s national CH<sub>4</sub> emissions in 2022 was 30 Gg CH<sub>4</sub>, so Georgia’s 2022 emissions were estimated to be (30 Gg CH<sub>4</sub> × 8.59% = 2.6 Gg CH<sub>4</sub>).

**Table 6-4. Pulp and Paper Permits Manually Removed From or Added to Analysis**

NPDES Stormwater/Construction or Non-Mill Permit Numbers Removed	NPDES Permit Numbers Added (Consistent with previous Inventories)
ALG060506, ALG060521, ALG141038, ARG160040, ARR001954, ARR00A499, ARR00A634, ARR00A771, ARR00A776, ARR156061, ARR157067, ARR157281, ARR157554, FLG071465, FLR05A517, FLR05B628, FLR05F649, FLR05G876, FLR05H761, FLR20CA56, FLR20CJ24, GAIS00705, GAIS00742, GAIS01355, GAIS01421, GAIS01678, GAIS02742, GAIS02751, GAIS02887, GAIS03756, GAIS04381, GAIS12083, GAIS13069, GAIS13100, GAIS13110, GAIS13795, GAIS14256, IDR053113, INRM01785, LAR05M630, LAR05P618, LAR10O640, LAR10O712, MANOE3652, MAR053165, MAR053218, MDR003165, MDR003388, MER05B433, MER05B451, MER05B451, MER05B608, MER05B983, MER05B984, MER05C163, MER05C172, MER05C178, MER05C178, MER05C200, MER05C269, MER05C269, MERNEB567, MERNEB567, MI0001210, MIS111133, MIS210982, MSR000044, MSR000382, MSR001256, MSR002038, MSR107486, MSR110045, MSR110077, MSR110077, MSR110099, MSR110099, MSR110118, MSR110118, MSR321403, MSR321403, NCS000101, NCS000105, NCS000106, NCS000106, NCS000211, NCS000211, NHG360002, NHNOEJ036, NHR053059, NHR053105, NJG224901, NMNOE3331, NYR00A955, NYR00B038, NYR00B199, NYR00B504, NYR00C573, NYR00E290, NYR00F582, NYR00F629, NYR00F629, NYR00G104, NYR00G109, ORR109393, ORR10A484, ORR10A546, ORR10E944, ORR10F683, ORR10F683, ORR10F806, ORR10F806, ORR220121, ORR221181, ORR240152, ORR240152, ORR240316, ORR241051, ORR241051, ORR241118, ORR241300, ORR241300, ORR241524, ORR241524, SCR004687, SCR005224, SCR006095, SCR006186, SCR006186, TNR050417, TNR050850, TNR050850, TNR051032, TNR051032, TNR052093, TNR053851, TNR054024, TNR054024, TNR055894, TNR056437, TNR058104, TXR05DP51, TXR1505FH, TXR15193U, TXR1547FX, TXR15691U, TXR1569FF, VAN030049, VAN030133, VAN040066, VAN040070, VAN040073, VAR052485	GA0001988, KYR003292, MN0001431, MN0001643



### 6.2.1.3.2. Meat and Poultry Processing

- Annual U.S. and state-level production data for red meat processing and poultry processing data are available from USDA-NASS (as shown in Appendix F, Table F-8). Depending on the commodity, limited state-level data are available. Typically, USDA reports only break out the primary states where the commodity is processed and then present production in “other states.”
- For red meat processing:
  - EPA gathered state-level 2022 and 1990 average live weight and total head slaughtered for the following commodities: beef, calves, hogs, and lamb/mutton (USDA 2023a, 1991a). EPA retained 2021 and 2012 data from the 1990–2021 state-level production data, 2019 data from the 1990–2019 state-level production data, and 2020 and 2004 data from the 1990–2020 state-level production data.
    - U.S. territories and the District of Columbia are not included in USDA-reported data.
  - For total head slaughtered (thousand head):
    - To populate states for which specific production data are not disclosed by USDA (“D” states), EPA evenly divided the difference between the sum of the state-level data and the reported national-level total to those D states.
    - Similarly, USDA provided a total for New England states that was evenly distributed to those states noted (Connecticut, Maine, Massachusetts, New Hampshire, Rhode Island, and Vermont).
    - USDA provided a combined total for Delaware and Maryland, which was evenly distributed between the two states.
  - For average live weight (pounds):
    - EPA used the average of available state-level data and the national average to determine the appropriate average live weight for the remaining states (D states). This calculated value was applied to all D states.
    - Similarly, the reported average live weight value for New England states was applied to those states.
    - USDA provided a combined total for Delaware and Maryland, which was evenly distributed between the two states.
  - As with the national *Inventory*, EPA determined live weight killed (LWK) by multiplying the average live weight by the total head/1,000 to get to million pounds LWK.
  - EPA added the disaggregated red meat processing data by state and divided the data by the reported national production to determine the proportion distributed to states. Because of the estimated nature of the calculated values, the total state-level LWK is estimated at about 95% of the national total, so the percentages were normalized to 100%.
- For poultry processing:
  - EPA gathered state-level 2022 and 1990 poultry live weight data. EPA retained 2021 and 2012 data from the 1990–2021 state-level production data, 2019 data from the 1990–2019 state-level production data, and 2020 and 2004 data from the 1990–2020 state-level production data. Only young chickens, or broilers, had state-level data available. Turkeys and mature chickens did not.
    - Young turkey data were available by state. EPA assumed that states with young turkeys would be representative of turkey processing production; therefore, young turkey data were used as a proxy for total turkeys (USDA 2023b, 1991b).

- Young chickens were used to represent mature chicken processing production by state (USDA 2023b, 1991b).
  - To populate D states for 2022, EPA evenly divided the difference between the sum of the state-level data and the reported national-level total to those D states.
  - To populate D states for 1990, EPA first proxied the reported D states for 2012 because the individual states for 1990 were not available or reported by USDA. This was done to encourage time series consistency and avoid showing states known to have poultry processing as having no emissions for the industry. EPA acknowledges this method could attribute minor emissions to states without poultry in 1990. Then, as with 2022, EPA evenly divided the difference between the sum of the state-level data and the reported national-level total to those D states.
  - For turkeys and mature chickens, the proportion of young turkeys and young chickens, respectively, was multiplied by the national-level value to determine the pounds of processing production per state.
  - Those values were added together and then divided by the total poultry (young chickens, mature chickens, turkeys) values to determine the proportion of poultry LWK for states.
- To calculate CH<sub>4</sub> emissions, EPA:
  - Multiplied national red meat plant CH<sub>4</sub> emissions by the percentage of U.S. total meat processing and added that to the national poultry plant CH<sub>4</sub> emissions multiplied by the percentage of U.S. total poultry processing by state.
  - Multiplied the 2004 (from the 1990–2020 inventory), 2012, 2019 (from the 1990–2019 inventory), 2020 (from the 1990–2020 inventory), 2021 (from the 1990–2021 inventory), and 2022 state-level proportion of U.S. meat and poultry BOD treated on-site by the national effluent CH<sub>4</sub> emissions from meat and poultry.
  - For 1991–2003, used linear interpolation of 1990 and 2004 state-level proportions, for 2005–2011, used linear interpolation of 2004 and 2012 state-level proportions, and for and 2013–2018, used the 2012 and 2019 proportions. Multiplied those values by the national effluent CH<sub>4</sub> emissions from meat and poultry.
  - Added plant and effluent emissions for total state-level emissions.
- To calculate N<sub>2</sub>O emissions, EPA:
  - Multiplied the 2004 (from the 1990–2020 inventory), 2012, 2019 (from the 1990–2019 inventory), 2020 (from the 1990–2020 inventory), 2021 (from the 1990–2021 inventory), and 2022 state-level proportion of U.S. total nitrogen in both
    - 1) aerobically treated meat and poultry wastewater by the N<sub>2</sub>O emissions from meat and poultry processing wastewater treatment for each year in the time series and
    - 2) discharged meat and poultry wastewater by the N<sub>2</sub>O emissions from meat and poultry processing wastewater treatment effluent for each year in the time series.
  - For 1991–2003, used linear interpolation of 1990 and 2003, for 2005–2011 and 2013–2018, EPA used linear interpolation of 2004 and 2012, and 2012 and 2019 state-level proportions, respectively. Multiplied those values by the national effluent N<sub>2</sub>O emissions from meat and poultry.
  - Added plant and effluent emissions for total state-level emissions.

### 6.2.1.3.3. Vegetables, Fruits, and Juices Processing

- Annual U.S. production data for vegetables, fruits, and juices processing are available from USDA. Depending on the commodity, state-level data are available (as shown in Appendix F, Table F-9). Typically, USDA reports only identify the primary states where the commodity is processed. For example, production data on broccoli are provided for California and “other states,” while production data on asparagus are provided for Michigan, Washington, and “other states.”
  - U.S. territories and the District of Columbia are not included in the USDA-reported data.
- EPA determined that the most recent year with complete state-level production values is 2017 because USDA suspended the reporting of some state-level production values in 2018 and more notably in 2019–2022.
- Previously, to better inform the time series, EPA investigated an earlier year, determined 2012 to be complete, and subsequently determined the state-level production values for 2012. EPA previously investigated and included 2004 during the 1990–2020 Inventory.
- For processing production data:
  - State-level data for potato processing were not available. Instead, EPA used state-level potato production (i.e., the production of potatoes grown not processed) as a proxy to determine the states to include (USDA 2014).
  - For other vegetables, EPA gathered data for asparagus, broccoli, carrots, cauliflower, sweet corn, cucumber (for pickles), lima beans, green peas, snap beans, spinach, and tomatoes (USDA 2015a). Where USDA reported data for “other states,” those data were distributed equally among the commodities. EPA added the production for these commodities to determine the percentage of the U.S. total for all “other vegetables,” which is the production value used in the national *Inventory* (not the individual commodities).
  - Processed apples, grapes used for wine, and citrus fruits were also determined at a state level. For apples, where USDA reported data for “other states,” those data were distributed equally (USDA 2015b, 2015c).
  - Noncitrus fruits are split out into separate commodities (e.g., blueberries, sweet cherries<sup>38</sup>); no state-level data are available for the aggregated “noncitrus fruit” category. Therefore, EPA gathered the state-level “utilized production” data for these separate commodities to determine the appropriate states and relative percentage of utilized production for noncitrus fruits (USDA 2015c).
  - Processed noncitrus fruit data are typically calculated in the national *Inventory* as utilized production minus fresh minus apples minus grapes for wine; however, because of the intensive nature of gathering data for the separate commodities, “utilized production” was used as a proxy for processed production data.
- To calculate emissions, EPA calculated the 2004, 2012, and 2017 percentage of U.S. total BOD by state and multiplied that by the national vegetables and fruits emissions for each year in the time series.

---

<sup>38</sup> EPA gathered 2004 and 2017 production for apricots; avocados (2012 values reported as “not available”); blueberries, cultivated blueberries (2004 only), and wild blueberries; boysenberries (2004 only); sweet and tart cherries; coffee (2017 only); cranberries; dates; loganberries (2004 only); nectarines; olives; papaya (2012 Hawaii crop reported as “not available”), including guavas and pineapples (Hawaii crops, 2004 only); peaches; pears; plums; prunes (combined with plums in 2004); raspberries; and strawberries.

- For 2005–2011 and 2013–2016, EPA determined state-level proportions by linear interpolation of 2004 and 2012, and 2012 and 2017 values, respectively. Proportions for 2018–2022 were assumed to be the same as 2017.

#### 6.2.1.3.4. *Petroleum Refining*

- Annual production data are available from EIA within the Department of Energy (EIA 2024a), as shown in Appendix F (Table F-10).
- Because state-level data may reveal confidential data, production data are aggregated by Petroleum Administration for Defense Districts (PADDs). Production data for the following PADDs and subdistricts are available:
  - PADD I (East Coast)
    - Subdistrict A (New England): Connecticut, Maine, Massachusetts, New Hampshire, Rhode Island, and Vermont
    - Subdistrict B (Central Atlantic): Delaware, District of Columbia, Maryland, New Jersey, New York, and Pennsylvania
    - Subdistrict C (Lower Atlantic): Florida, Georgia, North Carolina, South Carolina, Virginia, and West Virginia
  - PADD II (Midwest): Illinois, Indiana, Iowa, Kansas, Kentucky, Michigan, Minnesota, Missouri, Nebraska, North Dakota, South Dakota, Ohio, Oklahoma, Tennessee, and Wisconsin
  - PADD III (Gulf Coast): Alabama, Arkansas, Louisiana, Mississippi, New Mexico, and Texas
  - PADD IV (Rocky Mountain): Colorado, Idaho, Montana, Utah, and Wyoming
  - PADD V (West Coast): Alaska, Arizona, California, Hawaii, Nevada, Oregon, and Washington
- Operating capacity by state is available from EIA (2024b) for 1990–2022.
- EPA created state-level annual production data for each year of the time series (1990–2022) by dividing the annual production for each PADD subdistrict by the percentage of operating capacity each state provided in that year.
- Petroleum operating capacity values were not available for 1996 and 1998. These values were linearly interpolated.
- Example: 2022 California emissions
  - California data are included in PADD V.
  - PADD V has a total of 27 refineries with an operating capacity of 2,659,271 barrels.
  - California has a total of 15 refineries with an operating capacity of 1,749,871 barrels (or 65.8% of PADD V capacity).
  - PADD V produced 1,000,921 barrels in 2022.
  - Estimate California production as  $1,000,921 \text{ barrels} \times 65.8\% = 658,633 \text{ barrels}$ .
  - Calculate California's percentage of national production ( $658,633 \text{ barrels} / 7,079,773 \text{ barrels} = 9\%$ ).
  - Calculate California emissions as national emissions  $\times$  percentage of national production ( $4.4 \text{ Gg CH}_4 \times 9\% = 0.4 \text{ Gg CH}_4$ ).

#### 6.2.1.3.5. *Starch-based Ethanol Production*

- State-level ethanol production data are available from EIA's State Energy Data System (SEDS) (EIA 2023) (as shown in Appendix F, Table F-11).

- Fuel ethanol production data, including denaturant, in thousand barrels are available for 1960–2021 (EIA 2023).
- EPA checked the difference between SEDS national production and the reported production in the national *Inventory* and found small differences—on average, a 0.9% difference for the time series—further confirming SEDS is a good source of state-level production.
- Typically, the most recent year of data is used as a surrogate for the last year of available production data. For example, during the 1990–2022 *Inventory by State*, 2021 production values were used for 2022. This is due to the timing of when production data are released versus to publication of the *Inventory by State*.
- Calculated the percentage of national production by state for every year, using the production data noted above.
- Calculated the state-level emissions by multiplying national emissions by percentage production by state.
- Example: 2021 California emissions
  - 2021 California production value is 2,293 thousand barrels.
  - National production for 2021 is 375,517 thousand barrels.
  - California produced 1.2% of the national production in 2021.
  - Calculate 2021 California emissions as national emissions × percentage of national production (5.9 Gg CH<sub>4</sub> × 0.6% = 0.04 Gg CH<sub>4</sub>).

#### 6.2.1.3.6. *Breweries*

- Annual production data by state are available from the Alcohol and Tobacco Tax and Trade Bureau (TTB 2024) (as shown in Appendix F, Table F-12).
  - Quarterly state-level production data are available for 2015–2022. Annual, state-level taxable production values are available for 2008–2020. The quarterly state-level production values are preferred and were used for 2015–2022. Data for earlier years of the time series are still not available, so the calculated percentage of national production for 2008 was used for 1990–2007.
  - In cases where one or two quarters of data were not disclosed due to confidentiality reasons, EPA averaged the remaining quarters and added that average to the annual sum to estimate the total annual production for that state.
  - In cases where no quarterly state-level data were available for 2015–2020 (Delaware, Florida, Missouri, and New Hampshire), EPA used the annual taxable removals data to proportion the unaccounted-for production data (the difference between the U.S. total production and the sum of the state-level quarterly production data). For 2021 and 2022, EPA forecasted the available 2015–2020 data to estimate production data for those years, due to annual taxable removals data not being available.
  - Data are not available broken out between craft and noncraft production, so the approach assumes each state has the same distribution of craft and noncraft production as the national distribution.
- Calculated the percentage of national production by state.
- Calculated the state-level emissions by multiplying national emissions by percentage production by state.
- Example: 2019 California emissions
  - California production is 20,948,150 barrels.

- National production is 180,866,990 barrels.
- California produces 11.2% of national production.
- Calculate California emissions as national emissions × percentage of national production (5.2 Gg CH<sub>4</sub> × 11.2% = 0.581 Gg CH<sub>4</sub>).

#### 6.2.1.4 Recalculations

Recalculations discussed here are specific to state-level production or disaggregated data. To see impacts from updates to national-level data, see the recalculations discussion in Section 7.2 of the Waste chapter (Chapter 7) in the national *Inventories*, available online at [https://www.epa.gov/system/files/documents/2024-04/us-ghg-inventory-2024-chapter-7-waste\\_04-17-2024.pdf](https://www.epa.gov/system/files/documents/2024-04/us-ghg-inventory-2024-chapter-7-waste_04-17-2024.pdf).

EPA updated the domestic methodology to include state-level proportions of septic versus centralized treatment based on available data (U.S. Census Bureau 1990). These updates, in conjunction with the changes to the national *Inventories*,<sup>39</sup> resulted in changes for 1990–2017 for all state-level domestic CH<sub>4</sub> and N<sub>2</sub>O emission estimates.

Updates to the following state-level industrial production data, in conjunction with national-level updates, resulted in changes for the entire time series for every state-level total industrial CH<sub>4</sub> and N<sub>2</sub>O emission estimates:

- **Pulp and paper.** Including 2019, 2020, and 2021 flow estimates for all available state data due to an updated methodology to determine/download flow data from ECHO, affecting all years.
- **Meat and poultry processing.** Including 1990 production data, affecting 1990–2003.
- **Breweries.** Updating to use state-level production data (rather than taxable removals), affecting 2015–2021.

#### 6.2.1.5 Uncertainty

The overall uncertainty associated with the 2022 national estimates of CH<sub>4</sub> and N<sub>2</sub>O from wastewater treatment and discharge were calculated using the 2006 IPCC Guidelines Approach 2 methodology (IPCC 2006). As described further in Chapter 7 of the national *Inventories* (EPA 2024a), levels of uncertainty in the national estimates in 2022 were –29%/+33% for CH<sub>4</sub> and –36%/+192% for N<sub>2</sub>O. State-level estimates have a higher uncertainty due to apportioning the national emissions estimates to each state based solely on state population (for domestic) or state industry sector production (for industrial). This approach does not address state-level differences in the type of wastewater treatment systems in use or in the conditions of the state's receiving waterbodies. State-level emissions for the time series were estimated based on limited years of state-level data, which also results in higher uncertainty for the state estimates. These assumptions were required due to the general lack of readily available state- or regional-level data. For more details on national-level uncertainty, see the uncertainty discussion in Section 7.2 of the Waste chapter (Chapter 7) in the national *Inventories*, available online at [https://www.epa.gov/system/files/documents/2024-04/us-ghg-inventory-2024-chapter-7-waste\\_04-17-2024](https://www.epa.gov/system/files/documents/2024-04/us-ghg-inventory-2024-chapter-7-waste_04-17-2024).

#### 6.2.1.6 Planned Improvements

Generally, EPA plans to review feedback from reviews of the state-level inventory methods and assess potential to use data sets identified in comments to see if they provide comparable data for all states or

---

<sup>39</sup> See Section 7.2, page 7-52, of the national *Inventories*.

most states. The steps outlined below may inform the potential improvements for both domestic and industrial state-level emissions estimates. EPA plans to undertake the following assessments as resources allow:

- Determine state-level sources for the type of wastewater treatment systems in use for municipal or domestic or for industrial wastewater (by industrial sector).
- Determine state-level sources for BOD or total nitrogen data in municipal or domestic wastewater or industrial wastewater (by industrial sector).
- As stated in Section 7.2 of the national *Inventory*, investigate additional sources for estimating wastewater volume discharged and discharge location for both domestic and industrial sources.

For individual industries, EPA notes the following potential improvements.

#### **6.2.1.6.1. Pulp and Paper Manufacturing**

- Investigate state-level sources for the production of pulp, paper, and paperboard.
- Investigate additional years of ECHO data to improve the time series. Part of this includes evaluating the facilities present year to year to confirm time series consistency.
- Investigate states where data are reported as not applicable and confirm emissions estimates do not apply. Pending findings, determine another source to estimate wastewater flow for these states.
- Refine criteria for evaluating stormwater permits and identifying duplicate permits that should be removed from analysis.

#### **6.2.1.6.2. Meat and Poultry Processing**

- Continue to investigate additional years of available USDA data for inclusion to improve the time series.
- Investigate the presence of meat and poultry processing in the U.S. territories or the District of Columbia and, pending findings, additional sources for estimating those emissions. For the District of Columbia, reach out to USDA-NASS to confirm if the District of Columbia is already included in reporting.

#### **6.2.1.6.3. Vegetables, Fruits, and Juices Processing**

- Continue to investigate other years of available USDA data for inclusion. EPA investigated 1990 and 2008 during the 1990–2022 *Inventory* and determined that those data sets are inconsistent with the current data sets.
- Investigate the presence of vegetables, fruits, and juices processing in the U.S. territories or the District of Columbia and, pending findings, additional sources for estimating those emissions. For the District of Columbia, reach out to USDA-NASS to confirm if the District of Columbia is already included in reporting.

#### **6.2.1.6.4. Starch-Based Ethanol Production**

- Investigate sources to break down wet and dry milling by state over the time series.

#### **6.2.1.6.5. Breweries**

Investigate sources to break down craft and noncraft breweries by state over the time series.

### 6.2.1.7 References

- EIA (U.S. Energy Information Administration) (2023) *State Energy Data System (SEDS): 1960–2021 (Complete)*. U.S. Department of Energy. Available online at: <https://www.eia.gov/state/seds/seds-data-complete.php>.
- EIA (2024a) *Refinery and Blender Net Production*. U.S. Department of Energy. Available online at: [https://www.eia.gov/dnav/pet/pet\\_pnp\\_refp\\_a\\_ep00\\_ypr\\_mbbl\\_a.htm](https://www.eia.gov/dnav/pet/pet_pnp_refp_a_ep00_ypr_mbbl_a.htm).
- EIA (2024b) *Number and Capacity of Petroleum Refineries*. U.S. Department of Energy. Available online at: <https://www.eia.gov/state/seds/seds-data-complete.php>.
- EPA (U.S. Environmental Protection Agency) (2024a) *Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990–2022*. EPA 430-R-24-004. Available online at: <https://www.epa.gov/ghgemissions/inventory-us-greenhouse-gas-emissions-and-sinks>.
- EPA (2024b) *Water Pollution Search*. Available online at: <https://echo.epa.gov/trends/loading-tool/water-pollution-search>.
- Instituto de Estadísticas de Puerto Rico (2021) *Estimados Anuales Poblacionales de los Municipios Desde 1950*. Accessed February 2021. Available online at: <https://censo.estadisticas.pr/EstimadosPoblacionales>.
- IPCC (Intergovernmental Panel on Climate Change) (2006) *2006 IPCC Guidelines for National Greenhouse Gas Inventories*. H.S. Eggleston, L. Buendia, K. Miwa, T. Ngara, and K. Tanabe (eds.). Institute for Global Environmental Strategies. Available online at: <https://www.ipcc-nggip.iges.or.jp/public/2006gl/>.
- NEBRA (North East Biosolids and Residuals Association) (2022) *U.S. National Biosolids Data*. Available online at: [https://static1.squarespace.com/static/601837d1c67bcc4e1b11862f/t/62f4f5fbae32804dd9f51ef6/1660220925356/National\\_BiosolidsDataSummary\\_NBDP\\_20220811.pdf](https://static1.squarespace.com/static/601837d1c67bcc4e1b11862f/t/62f4f5fbae32804dd9f51ef6/1660220925356/National_BiosolidsDataSummary_NBDP_20220811.pdf).
- TTB (Alcohol and Tobacco Tax and Trade Bureau) (2024) *Beer Statistics*. Available online at: <https://www.ttb.gov/beer/statistics>.
- U.S. Census Bureau (1990). *Historical Census of Housing Tables: Sewage Disposal, 1990*. Available online: [https://www2.census.gov/programs-surveys/decennial/tables/time-series/coh-sewage/sewage1990.txt?itid=lk\\_inline\\_enhanced-template](https://www2.census.gov/programs-surveys/decennial/tables/time-series/coh-sewage/sewage1990.txt?itid=lk_inline_enhanced-template)
- U.S. Census Bureau (2002) Table CO-EST2001-12-00. In: *Time Series of Intercensal State Population Estimates: April 1, 1990 to April 1, 2000*. Release date: April 11, 2002. Available online at: <https://www2.census.gov/programs-surveys/popest/tables/1990-2000/intercensal/st-co/co-est2001-12-00.pdf>.
- U.S. Census Bureau (2011) Table ST-EST00INT-01. In: *Intercensal Estimates of the Resident Population for the United States, Regions, States, and Puerto Rico: April 1, 2000 to July 1, 2010*. Release date: September 2011. Available online at: <https://www2.census.gov/programs-surveys/popest/datasets/2000-2010/intercensal/state/st-est00int-alldata.csv>.
- U.S. Census Bureau (2021a) Table NST-EST2020. In: *Annual Estimates of the Resident Population for the United States, Regions, States, and Puerto Rico: April 1, 2010 to July 1, 2020*. Release date: July 2021.
- U.S. Census Bureau (2021b) Table NST-EST2021-POP. In: *Annual Estimates of the Resident Population for the United States, Regions, States, District of Columbia, and Puerto Rico: April 1, 2020 to July 1, 2021*. Release date: December 2021.



- U.S. Census Bureau (2022) *International Database: World Population Estimates and Projections*. Accessed January 2022. Available online at: <https://www.census.gov/programs-surveys/international-programs/about/idb.html>.
- U.S. Census Bureau (2023) *International Database*. Accessed September 2023. Available online at: [https://www.census.gov/data-tools/demo/idb/#/trends?YR\\_ANIM=2020&dashPages=DASH&FIPS\\_SINGLE=US&COUNTRY\\_YEAR=2022&menu=trendsViz&TREND\\_RANGE=1990,2022&TREND\\_STEP=5&TREND\\_ADD\\_YRS=&FIPS=AQ,GQ,CQ,RQ,VQ&measures=POP&CCODE=AS,GU,MP,PR,US,VI&CCODE\\_SINGLE=US&COUNTRY\\_YR\\_ANIM=2022](https://www.census.gov/data-tools/demo/idb/#/trends?YR_ANIM=2020&dashPages=DASH&FIPS_SINGLE=US&COUNTRY_YEAR=2022&menu=trendsViz&TREND_RANGE=1990,2022&TREND_STEP=5&TREND_ADD_YRS=&FIPS=AQ,GQ,CQ,RQ,VQ&measures=POP&CCODE=AS,GU,MP,PR,US,VI&CCODE_SINGLE=US&COUNTRY_YR_ANIM=2022).
- USDA (U.S. Department of Agriculture) (1991a) *Livestock Slaughter: 1990 Summary*. Available online at: <https://downloads.usda.library.cornell.edu/usda-esmis/files/r207tp32d/00000397q/bn999b62f/LiveSlauSu-03-00-1991.pdf>.
- USDA (1991b) *Poultry Slaughter*. Available online at: <https://downloads.usda.library.cornell.edu/usda-esmis/files/3197xm04j/x920fz52j/cj82k9125/PoulSlau-07-06-1990.pdf>.
- USDA (2014) *Potatoes: 2013 Summary*. Available online at: <https://downloads.usda.library.cornell.edu/usda-esmis/files/fx719m44h/h128nh39g/g445cg71m/Pota-09-18-2014.pdf>.
- USDA (2015a) *Vegetables: 2014 Summary*. Available online at: <https://downloads.usda.library.cornell.edu/usda-esmis/files/02870v86p/b2773z217/br86b639p/VegeSumm-01-29-2015.pdf>.
- USDA (2015b) *Citrus Fruits: 2015 Summary*. Available online at: <https://downloads.usda.library.cornell.edu/usda-esmis/files/j9602060k/gt54kq48n/xk81jn69p/CitrFru-09-17-2015.pdf>.
- USDA (2015c) *Noncitrus Fruits and Nuts: 2014 Summary*. Available online at: <https://downloads.usda.library.cornell.edu/usda-esmis/files/zs25x846c/6108vd86r/3r074x570/NoncFruNu-07-17-2015.pdf>.
- USDA (U.S. Department of Agriculture) (2023a) *Livestock Slaughter: 2022 Summary*. Available online at: <https://downloads.usda.library.cornell.edu/usda-esmis/files/r207tp32d/8p58qs65g/g445dv089/lsan0423.pdf>
- USDA (U.S. Department of Agriculture) (2023b) *Poultry Slaughter: 2022 Summary*. Available online at: <https://downloads.usda.library.cornell.edu/usda-esmis/files/pg15bd88s/m613p944x/ht24xx05j/pslaan23.pdf>.
- Washington Department of Ecology (2022) *Discharge Monitoring Reports (DMR) Data*. Available online at: <https://apps.ecology.wa.gov/paris/DischargeMonitoringData.aspx>.
- Washington Department of Ecology (2024) *Facility Summary: Nippon Paper Industries USA Co.* Available online at: <https://apps.ecology.wa.gov/paris/FacilitySummary.aspx?FacilityId=18>.